

Differential Rate Laws vs. Integrated Rate Laws

- Differential rate law: *Rate* vs. concentration
- Integrated rate law: concentration vs. time

Integrated Rate Law for a First-Order Reaction

- For a reaction, $A \rightarrow \text{products}$, which is first-order in A , we can write the differential rate law

$$\text{Rate} = \frac{-d[A]}{dt} = k[A]$$

- Consider the change in concentration of A from its initial value $[A]_0$ to its value $[A]$ at some later time t . By integral calculus it can be shown

$$-\ln\left(\frac{[A]}{[A]_0}\right) = kt \quad \Rightarrow \quad \ln\left(\frac{[A]}{[A]_0}\right) = -kt$$

- L This expression is the *first-order integrated rate law* equation.
- U Another way of writing this expression:

$$\ln\left(\frac{[A]_0}{[A]}\right) = kt$$

Linear Form of the First-Order Integrate Rate Law


$$\ln\left(\frac{[A]}{[A]_o}\right) = -kt$$

- Expanding the logarithm and rearranging:

$$\ln[A] = -kt + \ln[A]_o$$

- This is a straight-line equation:

$$y = mx + b$$



Half-Life of a First-Order Reaction

- L For any reaction we define the *half-life* as that time required for the concentrations of a reactant to diminish to half of what its value was at some initial time.

$$\text{If } t = t_{1/2} \text{ Y } [A] = \frac{1}{2}[A]_0.$$

Relationship Between $t_{1/2}$ and k

Substituting into the first-order integrated rate law expression at time $t_{1/2}$, when $[A] = \frac{1}{2}[A]_0$:

$$\ln\left(\frac{[A]_0}{[A]}\right) = \ln\left(\frac{[A]_0}{\frac{1}{2}[A]_0}\right) = kt_{1/2}$$


$$\ln 2 = kt_{1/2}$$

$$0.693 = kt_{1/2}$$

$$t_{1/2} = \frac{0.693}{k}$$

Concentration vs. Time in Half-Lives

- L After each $t_{1/2}$ period, there is half as much reactant as what was present at the beginning of the half-life period.



Concentration vs. Time in Half-Life Periods General Equation for a First-Order Reaction

Let h be the number of half-life periods elapsed:

$$h = \frac{t}{t_{1/2}}$$

Then, at time t the concentration of A as a function of its initial concentration $[A]_0$ is

$$[A] = [A]_0 \left(\frac{1}{2}\right)^h$$

L The value of h does not have to be an integer. 7

Nuclear Decay

- L Radioactive decay is a first-order process.

$$\text{Rate} = A = \lambda N$$

where A = decay rate (activity)
 λ = decay constant (analogous to k)
 N = number of radioactive atoms

- L The following equations can be defined in analogy with the general integrated rate equation and half-life equations for a first-order reaction:

$$\ln (N_t/N_0) = -\lambda t$$

$$t_{1/2} = 0.693/\lambda$$

$$N_t = N_0(1/2)^{h}$$

$$A_t = A_0(1/2)^{h}$$

Important Terms and Equations for First-Order Reactions

Terms

$[A]$ = concentration of A at some elapsed time, t

$[A]_0$ = concentration of A at an initial point in the reaction, at time $t = 0$

k = rate constant for the reaction, in units of $(\text{time})^{-1}$

$t_{1/2}$ = half-life period; time required for the concentration of A to fall to half the value it had at the beginning of the half-life period; i.e. time it takes for $[A] = \frac{1}{2}[A]_0$

h = number of half-life periods elapsed = $t/t_{1/2}$

Important Terms and Equations for First-Order Reactions

Equations

Differential rate law (*Rate* vs. concentration):

$$\text{Rate} = k[\text{A}]$$

Integrated rate law (concentration vs. time):

$$\ln \frac{[\text{A}]_0}{[\text{A}]} = kt \qquad \ln \frac{[\text{A}]}{[\text{A}]_0} = -kt$$
$$\ln[\text{A}] = -kt + \ln[\text{A}]_0$$

A plot of $\ln[\text{A}]$ vs. time is a straight line whose slope is $-k$ and whose intercept is $\ln[\text{A}]_0$.

Half-life:

$$[\text{A}] = [\text{A}]_0 \left(\frac{1}{2}\right)^h$$

$$t_{1/2} = 0.693/k$$

Important Terms and Equations for Radioactive Decay

Terms

N_t = number of radioactive atoms at some elapsed time, t

N_0 = number of radioactive atoms at an initial point, at time $t = 0$

λ = decay constant in units of $(\text{time})^{-1}$ (analogous to k)

A = decay activity in units of disintegrations per time per gram (analogous to *Rate*)

$t_{1/2}$ = half-life period; time required for N_t to fall to half the value it had at the beginning of the half-life period; i.e. time it takes for $N_t = \frac{1}{2}N_0$

h = number of half-life periods elapsed = $t/t_{1/2}$

Important Terms and Equations for Radioactive Decay

Equations

Radioactive decay law (analogous to differential rate law):

$$A = \lambda N$$

Time dependence of N (analogous to integrated rate law):

$$\ln \frac{N_0}{N_t} = \lambda t \qquad \ln \frac{N_t}{N_0} = -\lambda t$$
$$\ln N_t = -\lambda t + \ln N_0$$

A plot of $\ln N_t$ vs. time is a straight line whose slope is $-\lambda$ and whose intercept is $\ln N_0$.

Half-life:

$$N_t = N_0 \left(\frac{1}{2}\right)^h \qquad \text{and} \qquad A_t = A_0 \left(\frac{1}{2}\right)^h$$

$$t_{1/2} = 0.693/\lambda$$