


**Concentration Change with Time
for a Reversible Reaction**
(One-Step Mechanism in Both Directions)



L At t_{eq} , when equilibrium is established,

$$\text{Rate} = \frac{-1}{2} \frac{d[\text{NO}_2]}{dt} = \frac{d[\text{N}_2\text{O}_4]}{dt} = 0$$

Kinetics of $2\text{NO}_2(\text{g}) \rightleftharpoons \text{N}_2\text{O}_4(\text{g})$

- From the Law of Mass Action, for the forward reaction

$$rate_f = k_f [\text{NO}_2]^2$$

and for the reverse reaction

$$rate_r = k_r [\text{N}_2\text{O}_4]$$

- At any point in the reaction, the overall rate, *Rate*, is

$$Rate = rate_f - rate_r = k_f [\text{NO}_2]^2 - k_r [\text{N}_2\text{O}_4]$$

- At equilibrium,

$$Rate = k_f [\text{NO}_2]^2 - k_r [\text{N}_2\text{O}_4] = 0$$

which means

$$k_f [\text{NO}_2]^2 = k_r [\text{N}_2\text{O}_4]$$

or

$$rate_f = rate_r$$

- L This defines a state of *dynamic equilibrium*.

Defining the Equilibrium Constant for $2\text{NO}_2(\text{g}) \rightleftharpoons \text{N}_2\text{O}_4(\text{g})$

- At equilibrium

$$k_f[\text{NO}_2]^2 = k_r[\text{N}_2\text{O}_4]$$

- Rearranging to put the constants on one side

$$\frac{k_f}{k_r} = \frac{[\text{N}_2\text{O}_4]}{[\text{NO}_2]^2}$$

- Defining $K_c = k_f/k_r$

$$K_c = \frac{[\text{N}_2\text{O}_4]}{[\text{NO}_2]^2}$$


- L K_c is the **equilibrium constant**, defined as the ratio of product *equilibrium* concentration raised to its stoichiometric power over reactant *equilibrium* concentration raised to its stoichiometric power.
- L K_c is a constant for the reaction at a particular temperature, regardless of the starting concentrations.

K_c for a Multi-Step Mechanism



$$\text{Rate} = k[\text{NO}_2\text{Cl}]$$

- This is believed to proceed by the following two-step mechanism:



- Principle of Microscopic Reversibility:** If the reaction is at equilibrium overall, each elementary step and its reverse in the mechanism must also be at equilibrium.

$$K_1 = \frac{k_1}{k_{-1}} = \frac{[\text{NO}_2][\text{Cl}]}{[\text{NO}_2\text{Cl}]}$$
$$K_2 = \frac{k_2}{k_{-2}} = \frac{[\text{NO}_2][\text{Cl}_2]}{[\text{NO}_2\text{Cl}][\text{Cl}]}$$

- Multiplying $K_1 \times K_2$:

$$K_1 \times K_2 = \frac{k_1 k_2}{k_{-1} k_{-2}} = \frac{[\text{NO}_2][\text{Cl}]}{[\text{NO}_2\text{Cl}]} \frac{[\text{NO}_2][\text{Cl}_2]}{[\text{NO}_2\text{Cl}][\text{Cl}]} = \frac{[\text{NO}_2]^2[\text{Cl}_2]}{[\text{NO}_2\text{Cl}]^2}$$
$$= K_c$$

General Form of the Equilibrium Constant, K_c

L For a general reaction of the form




regardless of the number of steps in the mechanism, at equilibrium the concentrations of all species will have specific values that define a constant of the form

$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}$$


Rules for Writing K_c Expressions

1. For K_c , all concentrations are the values *at equilibrium* in mol/L.
2. All solids and neat liquids (not solutions) behave as if they have unit concentration and do not appear in the expression for K_c .
3. For solutions, solutes appear in the expression for K_c , but solvents are usually omitted unless active participants in the reaction.
4. Non-reactive species, such as spectator ions, are omitted from K_c .
5. K_c is defined for the reaction proceeding in the usual left-to-right manner, as written. The value of K_c for the reverse reaction is the inverse of that for the forward reaction.


Examples of K_c



$$K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$



$$K_c = \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]}$$



$$K_c = \frac{1}{[\text{CO}_2]}$$



$$K_c = \frac{[\text{H}_3\text{O}^+][\text{F}^-]}{[\text{HF}]}$$