

Review of Matrices

- L A matrix is a rectangular array of numbers that combines with other such arrays according to specific rules.

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdots & \cdot \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- T The dimension of a matrix is given as rows x columns; i.e., $m \times n$.

Matrix Multiplication

L If two matrices are to be multiplied together they must be *conformable*; i.e., the number of columns in the first (left) matrix must be the same as the number of rows in the second (right) matrix.

T The product matrix has as many rows as the first matrix and as many columns as the second matrix.


$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{bmatrix} = \begin{bmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{bmatrix}$$

T The elements of the product matrix, c_{ij} , are the sums of the products $a_{ik}b_{kj}$ for all values of k from 1 to m ; i.e.,

$$c_{ij} = \sum_{k=1}^m a_{ik}b_{kj}$$

$$\begin{bmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{bmatrix} = \begin{bmatrix} (a_{11}b_{11} + a_{12}b_{21}) & (a_{11}b_{12} + a_{12}b_{22}) & (a_{11}b_{13} + a_{12}b_{23}) \\ (a_{21}b_{11} + a_{22}b_{21}) & (a_{21}b_{12} + a_{22}b_{22}) & (a_{21}b_{13} + a_{22}b_{23}) \\ (a_{31}b_{11} + a_{32}b_{21}) & (a_{31}b_{12} + a_{32}b_{22}) & (a_{31}b_{13} + a_{32}b_{23}) \end{bmatrix}$$

Transformations of a General Vector in C_{2v}



$$E \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$C_2 \quad \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -x \\ -y \\ z \end{bmatrix}$$

$$F_v = F_{xz} \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x \\ -y \\ z \end{bmatrix}$$

$$F'_v = F_{yz} \quad \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -x \\ y \\ z \end{bmatrix}$$

A Representation with Matrices

| C_{2v} | E | C_2 | F_v | F_v' |
|------------|---|---|--|--|
| Γ_m | $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ | $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ | $\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ | $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ |

- T These matrices combine with each other in the same ways as the operations, so they form a representation of the group.
- T Γ_m is a **reducible representation**

A Representation from the Traces (Characters) of the Matrices

| | | | | |
|------------|-----|-------|-------|--------|
| C_{2v} | E | C_2 | F_v | F_v' |
| Γ_v | 3 | -1 | 1 | 1 |

T The characters of Γ_v are the sums of the corresponding characters of the three irreducible representations $A_1 + B_1 + B_2$:

| | | | | |
|------------|-----|-------|-------|--------|
| C_{2v} | E | C_2 | F_v | F_v' |
| A_1 | 1 | 1 | 1 | 1 |
| B_1 | 1 | -1 | 1 | -1 |
| B_2 | 1 | -1 | -1 | 1 |
| Γ_v | 3 | -1 | 1 | 1 |

○ $\Gamma_v = A_1 + B_1 + B_2$

T Breaking down Γ_v into its component irreducible representations is called **reduction**.

T The species into which Γ_v reduces are the those by which the vectors \mathbf{z} , \mathbf{x} , and \mathbf{y} transform, respectively.

Reduction of Γ_m by Block Diagonalization

| C_{2v} | E | | | C_2 | | | F_v | | | F_v' | | | |
|------------|-----|---|---|-------|----|---|-------|----|---|--------|---|---|-------|
| Γ_m | 1 | 0 | 0 | -1 | 0 | 0 | 1 | 0 | 0 | -1 | 0 | 0 | B_1 |
| | 0 | 1 | 0 | 0 | -1 | 0 | 0 | -1 | 0 | 0 | 1 | 0 | B_2 |
| | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 1 | A_1 |

- T Each diagonal element, c_{ii} , of each operator matrix expresses how one of the coordinates x , y , or z is transformed by the operation.
- Each c_{11} element expresses the transformation of the x coordinate.
 - Each c_{22} element expresses the transformation of the y coordinate.
 - Each c_{33} element expresses the transformation of the z coordinate.
- T The set of four c_{ii} elements with the same i (across a row) is an irreducible representation.
- T The three irreducible representations found by block diagonalization of Γ_m are the same as those found for Γ_v ; i.e.,

$$\Gamma_m = A_1 + B_1 + B_2 = \Gamma_v$$

Dimensions of Representations

- ⌊ In a representation of matrices, such as Γ_m , the *dimension of the representation* is the order of the square matrices of which it is composed.

$$d(\Gamma_m) = 3$$

- ⌊ For a representation of characters, such as Γ_v , the dimension is the value of the character for the identity operation.


$$P(E) = 3 \quad \text{Y} \quad d(\Gamma_v) = 3$$

- ⌊ The dimension of the reducible representation must equal the sum of the dimensions of all the irreducible representations of which it is composed.

$$d_r = \sum_i n_i d_i$$

More Complex Groups and Standard Character Tables

| C_{3v} | E | $2C_3$ | $3\sigma_v$ | | |
|----------|-----|--------|-------------|---------------------|-------------------------|
| A_1 | 1 | 1 | 1 | z | x^2+y^2, z^2 |
| A_2 | 1 | 1 | -1 | R_z | |
| E | 2 | -1 | 0 | $(x, y) (R_x, R_y)$ | $(x^2-y^2, xy)(xz, yz)$ |



- L The group C_{3v} has:
 - T Three classes of elements (symmetry operations).
 - T Three irreducible representations.
 - T One irreducible representation has a dimension of $d_i = 2$ (*doubly degenerate*).

- L The character table has a last column for *direct product transformations*.

Classes

- ⌊ **Geometrical Definition** (Symmetry Groups): Operations in the same class can be converted into one another by changing the axis system through application of some symmetry operation of the group.

- ⌊ **Mathematical Definition** (All Groups): The elements A and B belong to the same class if there is an element X within the group such that $X^{-1}AX = B$, where X^{-1} is the inverse of X (i.e., $XX^{-1} = X^{-1}X = E$).

- ⌋ If $X^{-1}AX = B$, we say that B is the *similarity transform* of A by X , or that A and B are *conjugate* to one another.

- ⌋ The element X may in some cases be the same as either A or B .

Classes of C_{3v} by Similarity Transforms

| C_{3v} | E | C_3 | C_3^2 | F_1 | F_2 | F_3 |
|----------|---------|---------|---------|---------|---------|---------|
| E | E | C_3 | C_3^2 | F_1 | F_2 | F_3 |
| C_3 | C_3 | C_3^2 | E | F_3 | F_1 | F_2 |
| C_3^2 | C_3^2 | E | C_3 | F_2 | F_3 | F_1 |
| F_1 | F_1 | F_2 | F_3 | E | C_3 | C_3^2 |
| F_2 | F_2 | F_3 | F_1 | C_3^2 | E | C_3 |
| F_3 | F_3 | F_1 | F_2 | C_3 | C_3^2 | E |

- L Take the similarity transforms on C_3 to find all members in its class:

$$EC_3E = C_3$$

$$C_3^2 C_3 C_3 = C_3^2 C_3^2 = C_3$$

$$C_3 C_3 C_3^2 = C_3 E = C_3$$

$$F_1 C_3 F_1 = F_1 F_3 = C_3^2$$

$$F_2 C_3 F_2 = F_2 F_1 = C_3^2$$

$$F_3 C_3 F_3 = F_3 F_2 = C_3^2$$

T Only C_3 and C_3^2

- L Take the similarity transforms on F_1 to find all members in its class:

$$E F_1 E = F_1$$

$$C_3^2 F_1 C_3 = C_3^2 F_2 = F_3$$

$$C_3 F_1 C_3^2 = C_3 F_3 = F_2$$


$$F_1 F_1 F_1 = F_1 E = F_1$$

$$F_2 F_1 F_2 = F_1 C_3 = F_2$$

$$F_3 F_1 F_3 = F_1 C_3^2 = F_3$$

T Only F_1 , F_2 , and F_3

Transformations of a General Vector in C_{3v} The Need for a Doubly Degenerate Representation



L No operation of C_{3v} changes the z coordinate.

T Every operation involves an equation of the form

$$\begin{bmatrix} ? & ? & 0 \\ ? & ? & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ z \end{bmatrix}$$

T We only need to describe any changes in the projection of \mathbf{v} in the xy plane.

L The operator matrix for each operation is generally unique, but all operations in the same class have the same character from their operator matrices.

T We only need to examine the effect of one operation in each class.

Transformations by E and $F_1 = F_{xz}$


E

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$F_1 = F_{xz}$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x \\ -y \\ z \end{bmatrix}$$

Transformation by C_3



From trigonometry:

$$x' = \cos \frac{2\pi}{3} x - \sin \frac{2\pi}{3} y = -\frac{1}{2}x - \frac{\sqrt{3}}{2}y$$

$$y' = \sin \frac{2\pi}{3} x + \cos \frac{2\pi}{3} y = \frac{\sqrt{3}}{2}x - \frac{1}{2}y$$

Therefore, the transformation matrix has nonzero off-diagonal elements:

$$\begin{bmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \left(-\frac{x}{2} - \frac{\sqrt{3}y}{2} \right) \\ \left(\frac{\sqrt{3}x}{2} - \frac{y}{2} \right) \\ z \end{bmatrix} = \begin{bmatrix} x' \\ y' \\ z' \end{bmatrix}$$

Reduction by Block Diagonalization

| C_{3v} | E | C_3 | | | F_v | | | |
|----------|-----|-------|------|------|-------|---|----|---|
| 1 | 0 | 0 | -1/2 | -1/2 | 0 | 1 | 0 | 0 |
| 0 | 1 | 0 | 1/2 | -1/2 | 0 | 0 | -1 | 0 |
| 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 1 |

- L The blocks must be the same size across all three matrices.

- T The presence of nonzero, off-diagonal elements in the transformation matrix for C_3 restricts us to diagonalization into a 2x2 block and a 1x1 block.

- T For all three matrices we must adopt a scheme of block diagonalization that yields one set of 2x2 matrices and another set of 1x1 matrices.

Representations of Characters

- L Converting to representations of characters gives a doubly degenerate irreducible representation and a nondegenerate representation.

| C_{3v} | E | $2C_3$ | $3C_2$ |
|---|-----|--------|--------|
| $\begin{matrix} x, y \\ = E \end{matrix}$ | 2 | -1 | 0 |
| $\begin{matrix} z \\ = A_1 \end{matrix}$ | 1 | 1 | 1 |

- T Any property that transforms as E in C_{3v} will have a companion, with which it is degenerate, that will be symmetrically and energetically equivalent.

| C_{3v} | E | $2C_3$ | $3C_2$ | | |
|----------|-----|--------|--------|---------------------|---------------------------|
| A_1 | 1 | 1 | 1 | z | $x^2 + y^2, z^2$ |
| A_2 | 1 | 1 | -1 | R_z | |
| E | 2 | -1 | 0 | $(x, y) (R_x, R_y)$ | $(x^2 - y^2, xy)(xz, yz)$ |

- T Unit vectors \mathbf{x} and \mathbf{y} are degenerate in C_{3v} .
- T Rotational vectors $\mathbf{R}_x, \mathbf{R}_y$ are degenerate in C_{3v} .

Direct Product Listings

| C_{3v} | E | $2C_3$ | $3C_2$ | | |
|----------|-----|--------|--------|---------------------|-------------------------|
| A_1 | 1 | 1 | 1 | z | x^2+y^2, z^2 |
| A_2 | 1 | 1 | -1 | R_z | |
| E | 2 | -1 | 0 | $(x, y) (R_x, R_y)$ | $(x^2-y^2, xy)(xz, yz)$ |

L The last column of typical character tables gives the transformation properties of direct products of vectors.

T Among other things, these can be associated with the transformation properties of d orbitals in the point group.

Correspond to d orbitals: $z^2, x^2-y^2, xy, xz, yz, 2z^2 - x^2 - y^2$

Do not correspond to d orbitals: $x^2, y^2, x^2 + y^2, x^2 + y^2 + z^2$

Complex-Conjugate Paired Irreducible Representations

- L Some groups have irreducible representations with imaginary characters in complex conjugate pairs:

$$C_n (n \geq 3), C_{nh} (n \geq 3), S_{2n}, T, T_h$$

- T The paired representations appear on successive lines in the character tables, joined by braces ($\{ \}$).
- T Each pair is given the single Mulliken symbol of a doubly degenerate representation (e.g., $E, E_1, E_2, E', E'', E_g, E_u$).
- T Each of the paired complex-conjugate representations is an irreducible representation in its own right.

Combining Complex-Conjugate Paired Representations

L It is sometimes convenient to add the two complex-conjugate representations to obtain a representation of real characters.

T When the pair has χ and χ^* characters, where $\chi = \exp(2Bi/n)$, the following identities are used in taking the sum:

$$\chi^p = \exp(2Bpi/n) = \cos 2Bp/n + i \sin 2Bp/n$$

$$\chi^{*p} = \exp(-2Bpi/n) = \cos 2Bp/n - i \sin 2Bp/n$$

which combine to give

$$\chi^p + \chi^{*p} = 2 \cos 2Bp/n$$

Example: In C_3 , $\chi = \exp(2Bi/3)$ and $\chi + \chi^* = 2 \cos 2B/3$.

| C_3 | E | C_3 | C_3^2 |
|---------|-----|---------------|---------------|
| E^a | 1 | χ | χ^* |
| E^b | 1 | χ^* | χ |
| $\{E\}$ | 2 | $2 \cos 2B/3$ | $2 \cos 2B/3$ |

L If complex-conjugate paired representations are combined in this way, realize that the real-number representation is a *reducible* representation.

Mulliken Symbols Irreducible Representation Symbols

In non-linear groups:

A nondegenerate; symmetric to C_n ($\chi_{C_n} > 0$)

B nondegenerate; antisymmetric to C_n ($\chi_{C_n} < 0$)

E doubly degenerate ($\chi_E = 2$)

T triply degenerate ($\chi_E = 3$)

G four-fold degenerate ($\chi_E = 4$) in groups I and I_h

H five-fold degenerate ($\chi_E = 5$) in groups I and I_h

In linear groups C_{4v} and D_{4h} :

E / A nondegenerate; symmetric to C_4 ($\chi_{C_\infty} = 1$)

$\{A, \sigma, M\} / E$ doubly degenerate ($\chi_E = 2$)

Mulliken Symbols Modifying Symbols

With any degeneracy in any centrosymmetric groups:

| | |
|---------------|---|
| subscript g | (<i>gerade</i>) symmetric with respect to inversion ($\chi_i > 0$) |
| subscript u | (<i>ungerade</i>) antisymmetric with respect to inversion ($\chi_i < 0$) |

With any degeneracy in non-centrosymmetric nonlinear groups:

| | |
|-------------------|---|
| prime (') | symmetric with respect to F_h ($\chi_{\sigma_h} > 0$) |
| double prime ('') | antisymmetric with respect to F_h ($\chi_{\sigma_h} < 0$) |

With nondegenerate representations in nonlinear groups:

| | |
|-------------|---|
| subscript 1 | symmetric with respect to C_m ($m < n$) or F_v ($\chi_{C_m} > 0$ or $\chi_{\sigma_v} > 0$) |
| subscript 2 | antisymmetric with respect to C_m ($m < n$) or F_v ($\chi_{C_m} < 0$ or $\chi_{\sigma_v} < 0$) |

With nondegenerate representations in linear groups (C_{4v} , D_{4h}):

| | |
|---------------|---|
| superscript + | symmetric with respect to $4F_v$ or $4C_2$ ($\chi_{\sigma_v} = 1$ or $\chi_{C_2} = 1$) |
| superscript - | antisymmetric with respect to $4F_v$ or $4C_2$ ($\chi_{\sigma_v} = -1$ or $\chi_{C_2} = -1$) |